

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231836015>

Zai improves nutrient and water productivity in the Ethiopian highlands

Article in *Experimental Agriculture* · January 2011

DOI: 10.1017/S0014479710000803

CITATIONS

37

READS

996

3 authors:



Tilahun Amede

Alliance for a Green Revolution in Africa

157 PUBLICATIONS 2,690 CITATIONS

[SEE PROFILE](#)



Mesfin Menza

Arba Minch University

11 PUBLICATIONS 87 CITATIONS

[SEE PROFILE](#)



Seleshi BEKELE Awlache

United Nations, New York

91 PUBLICATIONS 3,438 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Meta-analysis of crop response to fertilizer as a basis to design site-specific fertilizer recommendation [View project](#)



Integrated Watershed Management [View project](#)

ZAI IMPROVES NUTRIENT AND WATER PRODUCTIVITY IN THE ETHIOPIAN HIGHLANDS

By TILAHUN AMEDE†,‡, MESFIN MENZA§ and SELESHI BEKELE
AWLACHEW‡

†*International Livestock Research Institute and Challenge Programme on Water and Food,
P.O. Box 5689, Addis Ababa, Ethiopia*, ‡*International Water Management Institute,
Addis Ababa, Ethiopia* and §*Wollega University, Ethiopia*

(Accepted 7 August 2010)

SUMMARY

In the East African highlands, crop yields tend to increase with proximity of the farm plots to homesteads. Farmers identified soil erosion as the most detrimental cause of low crop yield in the outfields followed by soil compaction due to livestock trampling. The main objective of this study was to determine whether *zai* pits (i.e. small water harvesting pits) developed for dryland regions of the Sahel could increase crop yield and water productivity of degraded outfields in high rainfall areas, where mean annual rainfall exceeds 1300 mm but soil water infiltration is reduced by slope, low soil organic matter and hardpans. The pits were enlarged to resist strong runoff flows. The research was conducted over three years from 2004 to 2006. Potatoes and beans were used as test crops. Overall, compared to control plots, the *zai* pits, in combination with nitrogen (N) inputs, increased potato yields from 500% to 2000% ($p \leq 0.001$). The pits contributed more to increased crop yield than N inputs. Similarly, bean yields from the *zai* pits were up to 250% higher. Crop water productivity was 300–700% higher with *zai* pits than with control plots. The income of farmers who used *zai* pits was up to 20-fold higher than the labour costs required to prepare them. Contrary to conventional wisdom, this study reveals that the major constraint of the outfields is not nutrient deficiency *per se* rather low soil water holding capacity, which hinders crop growth and efficient utilization of available nutrients.

INTRODUCTION

Traditional land use in the Ethiopian highlands has failed to cope with increasing demand for food, feed and fibre. Land degradation, in combination with intermittent drought and declining soil fertility, is the major cause of low crop yield and food insecurity in the Ethiopian highlands. Estimates vary considerably but direct losses of productivity from land degradation are at least 3% of the agriculture gross domestic product of Ethiopia, while the loss of agricultural value during 2000–2010 was estimated to be \$7 billion (Berry, 2003). Exploitative traditional land use, aggravated by erosion, poor land management and increasing population, commonly led to nutrient mining and poor soil water holding capacity, particularly in fields far away from homesteads (Amede and Taboge, 2007). Moreover, very low levels of agricultural inputs, shortage of labour for agricultural production and conservation, and competing demands on crop and vegetative biomass for food, animal feed and soil fertility replenishment substantially reduced land and water productivity (Amede *et al.*, 2009).

†Corresponding author: T.Amede@cgiar.org

In the southern Ethiopian highlands, farmers tend to deliberately create fertile spots around homesteads. This is particularly pronounced in farming systems where food security is dependent on perennial food and cash crops such as banana and coffee and where theft and free grazing are problematic. The homestead plots are favoured for application of household refuse, manure and night soils, but are also enriched by nutrients coming from the outfields in the form of feed and mulch (Amede *et al.*, 2001). This results in a soil fertility gradient from homesteads to the outfields across farms and landscape (Amede and Taboge, 2007; Elias *et al.*, 1998). Accordingly, crop yields are higher in the plots closest to the houses. For instance, enset (*Enset ventricosum*) yields in outfields have been found to be 70% lower than those grown in homesteads (Amede and Taboge, 2007), and in a study from the same site, Elias *et al.* (1998) found that there was a positive nitrogen (N) and phosphorus (P) balance in the homesteads, compared to a strong negative balance in the outfields. The same trend was observed in a banana-based system in Uganda (Bekunda, 1999) and in western Kenya (Vanlauwe *et al.*, 2007). However, soil fertility decline and the associated decrease in crop yield in the outfields could not be explained solely by nutrient deficiency, since outfields rarely respond to fertilizer applications (Vanlauwe *et al.*, 2007) and crop yields do not also necessarily correlate with plant nutrient contents (Amede and Taboge, 2007).

In the same way that soil fertility declines, soil water content decreases from homesteads to the outfields, and spatial differences in soil water availability related to slope positions contribute to variable millet yields (Rockstrom and de Rouw, 1997). Despite relatively high average rainfall in the research area (1350 mm) and a seven-month growing period, crop water productivity (crop yield per unit of water depleted) is extremely low.

Despite lower productivity, outfields in the Ethiopian Highlands remain the largest arable land area for most farmers and they account for the largest share of total production (Elias *et al.*, 1998). Hence, any attempt to increase household food production and income in this region should focus on interventions that increase outfield productivity. Water productivity of these farms could be substantially increased by improving soil water infiltration and integrated water and nutrient management.

In this study, we tested the effects of small water harvesting pits, known as *zai* pits, which are commonly reported to increase crop yields in dry regions where low rainfall limits productivity (Abayomi *et al.*, 2001; Roose *et al.*, 1999). *Zai* pits are traditional practices developed in Burkina Fasso and commonly used in the Sudano-Sahelian areas for rehabilitating eroded and completely crusted fields, where the infiltration is too low to sustain vegetation (Roose *et al.*, 1999). *Zai* pits typically comprise holes dug during the dry season and then filled with one or two handfuls of dry dung (corresponding to 1–3 t ha⁻¹ of dry organic matter). These are then seeded with approximately a dozen cereal seeds after the first storms. The pits combine water harvesting and targeted application of organic amendments (Fatondji *et al.*, 2007). They have been shown to alleviate the adverse effects of dry spells (Hassen, 1996). In these studies, *zai* pits have been found to result in a 3- to 4-fold increases in grain yield of millet compared to flat planting, mainly as a consequence of *in situ* water conservation




Figure 1. Location map of the research sites in Gununo, Ethiopia.

(Fatondji *et al.*, 2007). The innovative feature of our study was to ascertain the effect of *zai* in a region with relatively high rainfall and sloping landscapes.

The principal objectives of this study were to: (i) evaluate the effect of *zai* pits along with fertilizer application on crop yield of potatoes and beans; (ii) evaluate the effects of these interventions on crop water productivity; and (iii) assess farmers' perception of the cause of land degradation in the outfields and their response to *zai*.

MATERIALS AND METHODS

Characteristics of the study area

The research sites are located at Gegecho and Offa villages of Gununo, South-Western Ethiopian highlands ($37^{\circ}39'E$, $6^{\circ}56'N$) (Figure 1). This area is one of the most densely populated districts in the country, with more than 400 people per km^2 . Farm sizes typically range from 0.35 to 0.6 ha for a family of seven (Amede and Delve, 2008). Most farms are located on hillsides with slopes ranging from 5% to 23%. Households tend to be located up slopes with the outfields located close to the valley bottoms.

The area has a bimodal mean annual rainfall of 1300 mm, an average temperature of 19.5°C and an altitude range of 1880–1960 m asl. The small rainy season (*belg*)




Figure 2. Monthly rainfall during the growing season (March to October) in Gununo, Ethiopia.

extends from March to June while the main rainy season (*meher*) extends from July to the end of October (Figure 2). The months of July and August receive the highest rainfall and cause significant soil loss, mainly in the outfields. The dominant soils are deep Eutric Nitisols (Amede *et al.*, 2001). These soils are dominated by kaolinitic minerals, which are inherently low in N and P (Table 1).

Mixed crop-livestock systems dominate where farmers cultivate small quantities of a large number of crops including enset, cereals, root crops and vegetables in different combinations. Like most farming systems in the East African highlands, the homestead and midfields (i.e. fields located between the homestead and the outfields) are mostly occupied by enset, coffee, taro and sweet potato (*Ipomea batatas*). The outfields are typically used for growing cereals, sweet potato, potato (*Solanum tuberosum*) and beans (*Phaseolus vulgaris*). The major cereals (in terms of area and household consumption) are maize (*Zea mays*) and teff (*Eragrostis abyssinica*). Farmers also cultivate sorghum (*Sorghum bicolor*) and wheat (*Triticum aestivum*) to a limited extent. The most widespread cropping pattern in the outfields is to alternate maize in the *belg* with sweet potato or teff in the *meher* (Elias *et al.*, 1998). The first season crops are planted in March and harvested in June or early July, and the second season crops are planted in July or August and harvested in October or November.

Survey with farmers

After consultative meetings with the community to understand major constraints of farmers, formal and informal surveys were conducted to establish the perception of farmers on the causes and effects of soil fertility decline in the outfields and to learn their views about the new intervention, *zai*. A formal survey with two rounds of questionnaire use was conducted between June and October 2004 with 20 key informants who the community identified as being 'good farmers' and are farming slope fields. Interviews were conducted by trained research officers who spoke the

local language. The results from the first survey, conducted in mid August, produced local criteria for identifying degraded fields. During the second survey, conducted in the third week of October, 2005, we identified 8 out of 32 local indicators that farmers considered to be very important, using pair-wise analysis methods (Table 2).

Experimental designs

In consultation with the key informants three representative farms, belonging to the informants and representing degraded fields in the community were identified. The three farms (Farms A, B and C) faced each other across the V-shape valley, and were used to evaluate the effects of *zai* pits in combination with manure and different levels of N application on potato and beans. Farm B and C were neighbouring farms and belonged to brothers. The experimental plots, labelled as degraded outfields, were located about 60 m from the homestead fields, commonly bordering grazing land. On 20 February 2005, shortly before the beginning of the short rainy season, soil samples were collected from the homestead and outfields of the three farms (A, B, C) and analysed for nitrogen (N), phosphorus (P), cation exchange capacity (CEC) and organic matter (OM) using the nutrient analysis procedures described below.

About 1 kg of soil samples were collected per spot from above 20 cm soil depth, in five replicates per farm. Soil samples were oven dried to constant weight, then ground to pass through a 1.0 mm sieve and analysed for total NPK by Kjeldahl digestion with concentrated sulphuric acid (Anderson and Ingram, 1993). N, P, CEC and OM were measured in the laboratory of the International Livestock Research Institute using standard procedures. Soil water content was measured from the top 30 cm of the soil using gravimetric methods during a dry spell, three weeks after rain. The soil nutrient and water content nutrient from three representative farms is presented in Table 1.

N was reported to be the major limiting nutrient affecting crop production in the outfields of this area (Elias *et al.*, 1998) and therefore was included in the experiment. A randomized block design with four replications and nine plots per replication, with a plot size of 9 m² was used. The treatments were a combination of the following treatments: (i) *zai* or flat planting, (ii) with or without manure and (iii) 0, 30 or 60 kg N ha⁻¹. The control treatments were those without *zai* and without manure. There were 1.5 m wide gangways between replications. The experiment was protected with 2 m of border crops, which were also separated by 2 m of gangways.

In the Sahel, the *zai* consists of digging holes during the dry season (30 cm diameter, 15 cm deep) then filling them with one or two handfuls of dry dung (Roose *et al.*, 1993). In 2004, we set a preliminary experiment to evaluate the effects of *zai* pits, following the designs from the Sahel (Abayomi *et al.*, 2001; Roose *et al.*, 1999). However, the pits were filled by soil and sediments in one day of torrential rain (data not shown). Following this experience, we opted for *zai* pits with a larger diameter of 50 cm and depth 45 cm. We created wider and deeper *zai* pits to capture the runoff and minimize the erosion effects. About 1.5 kg of low quality stall manure, which is equivalent to about 5.8 t ha⁻¹ collected from a heap, was thoroughly mixed with the soil and returned to the pits. The same amount of manure was broadcast for the flat planting.

N was applied in the form of urea. Tri super phosphate (TSP) was applied to provide 30 kg ha⁻¹ of P across all treatments during planting, reflecting the low-input farming practices of the system. We also put two rows of soil bunds above the experimental plots to protect them from erosion.

In 2005, six seed tubers were planted in each *zai* pit, three weeks after the incorporation of the manure. This corresponded to a seed rate of about 1.1 t ha⁻¹. Equal number of tubers was planted in the flat fields fertilized with manure. Urea treatment was applied three weeks after planting. The field was kept weed free by hand. The plants were harvested following physiological maturity.

In 2006, the experiment was repeated with beans on the same old *zai* pits with the application of urea at a similar rate of 0, 30 and 60 kg N ha⁻¹ and a basal dressing with 30 kg TSP. A bean variety, Omo-95, was planted in the *zai* pits and flat plots at a seed rate of 50 kg ha⁻¹ on 16 March 2006. The beans were harvested on 23 June 2006. The soil nutrient and soil water content of *zai* and flat plots were measured following the methods described above.

One major concern of adoption of *zai* is the labour cost. Labour cost to prepare the *zai* pits was calculated after considering the number of pits per ha and the number of man days per pit. Labour cost and farm gate price in 2006 (7 birr day⁻¹ and 500 birr t⁻¹ potato, respectively), were used to calculate costs and benefits (in 2006 1 US\$ = 8.4 Ethiopian birr). All other costs of inputs were the same for the *zai* pits and the flat fields. In all cases, although the farmers provided their land and labour, the experiments were controlled by researchers.

Establishing water productivity

Crop water productivity (CWP) of the various treatments was calculated after simulation of climatic data using New LocClim (FAO, 2005) and CropWat model (FAO, 1998). The crop coefficient (K_c) was estimated as a ratio of crop evapotranspiration (ET_c, mm day⁻¹) and reference evapotranspiration (ET_o) (FAO, 1998). Water productivity of control and treatment plots was calculated by dividing grain or tuber yield per total amount of water used per respective crop and treatment. Dry weight of potato (22%) was used to calculate CWP.

Statistical differences among treatments were determined through analysis of variance with ranks and the Tukey test using Sigma stat (Jandel Scientific, 1994).

RESULTS

Soil fertility gradients

There was a significant difference in P, OM and soil water content (SWC) between homesteads and outfields of the respective farms. The P and organic matter contents of homestead plots were at least three times higher than the outfields (Table 1). Farm C had lower contents of N and OM than Farms A and B. In addition, SWC also varied across soil fertility gradients and farms. At the time of measurement, the SWC of the outfields was on average 5% less than in the homesteads (Table 1). Although

Table 1. Soil nutrient status and soil water content of homesteads and outfields farms in Gununo, Ethiopia ($n = 5$ samples from within the field).

Parameters	Homestead fields (mean)			Outfields (mean)			<i>p</i> -value
	Farm A	Farm B	Farm C	Farm A	Farm B	Farm C	
Total N (%)	0.24	0.22	0.18	0.15	0.18	0.09	<i>n.s.</i>
Available phosphorus (ppm)	7.49	7.36	8.42	0.70	0.75	0.78	***
CEC (meq 100 g ⁻¹)	23.19	25.67	27.83	18.69	18.97	25.72	<i>n.s.</i>
Organic matter (%)	6.0	4.99	5.11	2.61	2.00	1.67	***
Soil water content (%)	18.1	20.23	21.17	14.17	15.80	17.62	*

n.s. = not significant, * and *** significantly different at $p \leq 0.05$ and $p \leq 0.001$ between homestead and outfields.

Table 2. Farmers ranking of potential causes of land degradation of outfields, using pair wise comparison in Gununo, Ethiopia ($n = 20$).

	Erosion	Labour shortage	Livestock trampling	Crop rotation	Crop residue availability	Management skills	Manure availability	Other factors
Sum	134.0	71.0	80.00	64.0	59.0	35.0	20.0	68.0
Mean	6.7	3.5	4.00	3.2	2.9	1.7	1.0	3.4
<i>s.e.</i>	0.5	1.4	2.03	0.7	1.1	1.5	1.5	2.7
% total	25.2	13.4	15.1	12.1	11.1	6.6	3.8	12.8

there was a higher N content and CEC in homestead plots compared to outfields, the difference between these fields was not significant.

Farmer's perceptions

Out of the 32 potential causes of land degradation (data not presented), 8 were identified by the farmers to be the most important in facilitating outfield degradation. Based on the results of the pair-wise analysis of the most important eight constraints, the major causes of land degradation, according to priority order (from most important to least important) were: soil erosion, livestock trampling, labour shortage to construct soil conservation bunds and transport manure during the growing season, limited crop rotation in the outfields, and shortage of manure and other biomass to fertilize the outfields (Table 2). However, farmers recognized the detrimental effect of erosion on land quality as problem number one. Potentially, these factors have induced preferential management of homestead fields over outfields.

Effects of zai pits and N levels on crop yield

There was significant difference among the three farms in response to the *zai* pit treatments and fertilizer application ($p \leq 0.05$) (Figure 3). Farm A was much more productive than the other two farms, both in the *zai* pits and the controls. Potato yield was significantly higher in *zai* pits than those planted in flat fields. In fact, in Farms B and C, tuber yield of control plots was approaching zero and was not measurable because of stunted plants that failed to produce tubers. The yield advantage of the *zai* system was about 5-fold in Farm A while it was about 20-fold in Farm C. The effect




Figure 3. Effects of direct application or incorporation of biomass to a *zai* plot on tuber yield of potato, 2005. Error bars indicate standard error of difference between means ($n = 4$).

of different levels of N on tuber yield was pronounced, mainly through the *zai* pits. Significantly high potato tuber yield was obtained from the *zai* with 60 kg N ha⁻¹ followed by 30 kg N ha⁻¹ in Farms B and C. There was no significant difference in yield between 30 kg N ha⁻¹ and 60 kg N ha⁻¹ on Farm A (Figure 3). Similarly, bean yield from the *zai* pits was about 2.5 times greater than that from flat planting (Figure 4). The effect of N application on beans grain yield was significant in both *zai* and flat planting.

In addition to crop yield, soil moisture content of the *zai* pits was significantly higher ($p < 0.01$) than that of flat planting (Figure 5). The difference in some plots was up to 10%. However, we did not find differences in nutrient content (N, P and CEC) between *zai* pits and flat plots, with N content in the soil being less than 1% and organic matter being less than 1.8% in both treatments.

By growing potato in about 3925 pits ha⁻¹, the farm income produced by *zai* pits was up to 20-times higher than the labour costs required to prepare them (data not shown). Hence, it was economical to use the *zai* pits for cultivating potatoes on the degraded farms.




Figure 4. Effects of different levels of nitrogen on grain yields of beans with or without *zai* pits, 2006. Error bars indicate standard error of difference between means ($n = 4$).




Figure 5. Organic matter and soil water content within and outside *zai* pits, 2005. Error bars indicate *s.d.* ($n = 4$).

Effects of zai on crop water productivity (CWP)

There was a significant correlation between water productivity and crop yield ($R = 0.98$). However, the actual productivity varied with both crop type and treatment. Water productivity was the highest with potato, up to 13.2 kg of produce per cubic metre of water used. The highest increase in water productivity was obtained from




Figure 6a and 6b. Crop water productivity of potato (a) and beans (b) under control or *zai* treatments in Gununo, Ethiopia. ND indicates no data collected.

the *zai* pits, ranging from 260 to 620% compared to the control. The benefit of N application on CWP was in the range of 5–40% for potato and 135–265% for beans (Figure 6). The effect of N application on CWP was significantly higher than the

control ($p \leq 0.05$) both in potato and beans, with higher returns at a rate of 60 kg ha⁻¹. However, the large difference in water productivity between potato and beans under *zai* pits is also partly due to the higher moisture content of potato tubers than bean seeds.

DISCUSSION

In contrast to conventional wisdom, this study reveals that the major constraint of the outfields is not nutrient deficiency *per se* but rather low soil water holding capacity, which hinders crop growth and efficient utilization of available nutrients. Conventional wisdom suggests that crop water requirements could be easily satisfied in highland rain-fed systems where rainfall supply exceeds evapotranspiration demands. This wisdom, together with the nutrient balance studies of 1990s, forced researchers to believe that yield differences between homestead fields and outfields in the East African highlands was strongly associated with differences in soil nutrient status. Some of the most comprehensive studies on this subject emerged from Stoorvogel and Smaling (1990) for sub-Saharan countries (a continental study), Smaling *et al.* (1993) for the Kisii district of Kenya and Elias *et al.* (1998) for Southern Ethiopia. Stoorvogel and Smaling (1990) suggested that an analysis of nutrient inputs and outputs could indicate whether the soil fertility is being maintained, improved or degraded. The results of these and other studies suggested nutrient depletion to be a major problem of outfields in this region. These studies also suggested that degradation of the outfields was associated with limited financial capacity of farmers to buy chemical fertilizers to maintain soil fertility (Elias *et al.*, 1998; Smaling *et al.*, 1993). Similarly, Amede and Taboge (2007) in a recent study reported a 70% yield reduction in the local food crop enset when grown in outfields. However, the plant nutrient content of the enset plants grown in the outfields was much higher than plants grown in homesteads, indicating a reduced utilization of these plant nutrients (Amede and Taboge, 2007). Moreover, application of chemical fertilizers alone to these degraded farms did not increase crop yield to the expected levels (Figure 3). Vanlauwe *et al.* (2001) also reported that the return from chemical fertilizers in degraded plots was much lower than the return from fertile soils implying to an additional limiting factors affecting crop production in these systems.

Erosion was also an important negative contributor to the nutrient balance of the outfields in these areas (Amede *et al.*, 2001; Elias *et al.*, 1998) as also confirmed by local farmers (Table 1). It facilitates the removal of soil organic matter, reducing the water holding capacity of the soil substantially through reduced porosity, poor water infiltration and increased run-off (Bossio *et al.*, 2007). These factors contributed to the decline in soil quality of outfields, which became shallower, with very high soil density and low organic matter content to capture and store water in the root zone, particularly in farms on steep slopes. All of these soil parameters are indicative of water-stressed outfields.

While there was a significant difference in soil water content of *zai* and flat planted plots, we did not find differences in organic matter content within and outside the *zai* pits (Figure 5) because of the longer period required for soil organic matter

build-up. Crop water productivity was much higher in potato than in beans (Figure 6), indicating genetic differences in water use efficiency between these crop species. Though application of manure increased potato yield by about 50%, potato yield from *zai* was significantly higher (by a factor of 5) ($p \leq 0.001$) than from application of low quality manure combined with N in the form of urea. Water productivity has also tripled with *zai* interventions without any considerable change in nutrient status of the soils (data not shown). Hence, we consider that the yield gains from *zai* pits were strongly associated with improved soil water availability at the critical growth stages rather than increased nutrient supply. It also suggests that higher soil water content in the *zai* pits was due more to physical conservation of water by the pits than contributions made by the manure. *Zai* pits were also found to improve the water productivity of millets in the Sahel by a factor of 2 (Fatondji *et al.*, 2007). In the same way, experiments conducted with traditional *zai* system for restoring a degraded Entisol in Burkina Faso showed that the highest production of grain (11 times the control) and straw (5 times the control) in *zai* pits was achieved mainly with a combination of organic amendment and mineral fertilizers (Roose and Barthès, 2001). These added benefits are caused by direct interactions among conserved water, decomposing organic matter and fertilizer-N leading to improved synchrony between supply of and demand for available N (Vanlauwe *et al.*, 2001) and soil water.

The effect of N application on crop yield significantly increased in *zai* pits, particularly at higher application rates, possibly due to minimized washing-away effects of erosion. Although an increase in soil organic matter content through incorporation of green manures was possible (Buerkert *et al.*, 2000) the benefits could be undermined by the high annual soil loss, which is between 35 and 80 t ha⁻¹ for Gununo, depending on the season (SCRIP, 1996) unless soil and water conservation structures (e.g. *zai* pits) are adopted by farmers.

As farms in these areas contain heterogeneous fields with different soil fertility status (Table 1) and slopes, these soil and water management interventions should be targeted within this heterogeneity. Hence, the labour-intensive interventions like *zai* should target the most affected outfields, particularly where other management options fail to bring impact. These interventions were economically viable. Moreover, labour shortage is not a major constraint for the majority of these households (Amede and Delve, 2008), particularly since digging of the *zai* holes should be done during the dry period when farmers can invest their spare time (Fatondji *et al.*, 2007). However, these interventions may not be appropriate for every farm or system though it could target farm niches. For instance, the level of adoption of *zai* could be low in plain fields where runoff is minimal and soil water infiltration is relatively high.

CONCLUSION

Very small land holdings (<0.5 ha for a family of seven) and occasional drought are considered as the major causes of food insecurity in SSA. However, food insecurity in these farming systems is also associated with low productivity of outfields, which cover more than 65% of the total crop land. Results from this work suggested that

household food production could be substantially increased by improved trapping of water and nutrients using *zai* pits in these outfields. They could be used to capture water and increase recharge of upslope and degraded watersheds. These interventions are equally effective in high rainfall, high slope areas, where soil water infiltration is reduced by slope, soil organic matter is low and hardpans are created due to erosion and improper livestock management. However, the scarcity of manure presents a constraint to the use of the *zai* technology unless biomass production for organic fertilization is promoted as an important intervention.

Acknowledgments We would like to thank Drs Don Peden and Matthew MacCartney for their valuable comments on the manuscript and the farmers for their committed engagement. The data was generated while the first author was staff of the African Highlands Initiative and TSBF-CIAT. We acknowledged the support of the staff of Areka research centre during data collection.

REFERENCES

- Abayomi, Y. A., Fadayomi, O., Babatola, J. O. and Tian, G. (2001). Evaluation of selected legume cover crops for biomass production, dry season survival and soil fertility improvement in a moist savannah location in Nigeria. *African Crop Science Journal* 9:615–627.
- Amede, T. and Delve, R. (2008). Modelling crop-livestock systems for achieving food security and increasing production efficiencies in the Ethiopian highlands. *Experimental Agriculture* 44:441–452.
- Amede, T. and Taboge, E. (2007). Enhancing farmer innovation through manipulation of soil fertility gradients in enset systems. In *Improving human welfare and environmental conservation by empowering farmers to combat soil fertility degradation*, 289–297. (Ed. A. Bationo). African Soils Network, Springer Verlag.
- Amede, T., Belachew, T. and Geta, E. (2001). Reversing the degradation of arable land in Ethiopian Highlands. *Managing African Soils No. 23*. International Institute for Environment and Development, London.
- Amede, T., Descheemaeker, K., Peden, D. and van Rooyen, A. (2009). Harnessing benefits from improved livestock water productivity in crop-livestock systems of sub-Saharan Africa: synthesis. *The Rangeland Journal* 31: 169–178.
- Anderson, J.M., and Ingram, J. S. L. (1993). *Tropical Soil biology and Fertility: A Handbook of Methods*. Wallingford, UK: CAB International.
- Bekunda, M. (1999). Farmers' response to soil fertility decline in banana-based cropping systems of Uganda. *Managing African Soils No. 4*. International Institute for Environment and Development, London.
- Berry, L. (2003). Land degradation in Ethiopia: Its extent and impact. *Commissioned paper by the GM and WP support*. Available from ftp://ftp.fao.org/agl/agll/ladadocs/ETHIOPIA_LD_CASE_STUDIES.doc
- Bossio, D., William, C. W., Geheb, K., Van Lynden, G. and Mati, B. (2007). Conserving land and protecting water. In *Water for Food, Water for Life: A Comprehensive Assessment of Water Management in Agriculture*, 551–583 (Ed. D. Molden) London: Earthscan.
- Buerkert, A., Bationo, A. and Dossa, K. (2000). Mechanisms of residue mulch-induced cereal growth increases in West Africa. *Soil Science Society of America. Journal* 64:346–358.
- Elias, S., Morse, S. and Belshaw, D. G. R. (1998). Nitrogen and phosphorus balances of Kindo Koisha farms in Southern Ethiopia. *Agriculture, Ecosystems and Environment* 71:93–113.
- Fatondji, D., Martius, C., Bielders, C. L., Vlek, P. L. G., Bationo, A. and Gerard, B. (2007). Effect of planting technique and amendment type on pearl millet yield, nutrient uptake and water use on degraded land in Niger. In *Improving Human Welfare and Environmental Conservation by Empowering Farmers to Combat Soil Fertility Degradation*, 179–193 (Ed. A. Bationo). African Soils Network, Springer Verlag.
- FAO (1998). *Crop evapotranspiration – guidelines for computing crop water requirements*. FAO, Rome.
- FAO (2005). *Local climate estimator* (New LockClim 1.06). FAO, Rome.
- Hassen, A. (1996). Improved traditional planting pits in the Tahoua department, Niger. An example of rapid adoption by farmers. In *Sustaining the soil. Indigenous soil and Water Conservation in Africa*, 56–61 (Eds C. Reij, I. Scoones and C. Toulmin). London: Earthscan.

- Jandel Scientific Software (1994). *Sigma Stat Statistical Software for Windows*. Jandel, San Rafael, CA.
- Rockstrom, J. and de Rouw, A. (1997). Water, nutrients and slope position in on-farm pearl millet cultivation in the Sahel. *Plant and Soil* 195: 311–327.
- Roose, E. and Barthès, B. (2001). Organic matter management for soil conservation and productivity restoration in Africa: a contribution from Francophone research. *Nutrient Cycling in Agroecosystems* 61:1–2.
- Roose, E., Kabore, V. and Guenat, C. (1999). *Zai* practice: a West African traditional rehabilitation system for semiarid degraded lands, a case study in Burkina Faso. *Arid Soil Research and Rehabilitation* 13: 343–355.
- Smaling, E. M. A., Stoorvogel, J. J. and Windmeijer, P. N. (1993). Calculating soil nutrient balances in Africa at different scales II. District scale. *Fertility Research* 35: 237–250.
- Soil Conservation Research Program (SCRIP) (1996). *Data Base Report (1982–1993)*, Series II: Gununo Research Unit. University of Berne, Berne.
- Stoorvogel, J. J. and Smaling, E. M. A. (1990). Assessment of soil nutrient depletion in sub-Saharan Africa 1983–2000. *Report 28. The Winand Staring Centre for Integrated Land, Soil and Water Research (SC-DLO), Wageningen*.
- Vanlauwe, B., Tittonell, P. and Mukalama, J. (2007). Within soil fertility gradients affect response of maize to fertilizer application in western Kenya. *Nutrient Cycling in Agroecosystems* 76:2–3.
- Vanlauwe, B., Aihou, K., Houngnandan, P., Diels, J., Sanginga, N. and Merckx, R. (2001). Nitrogen management in ‘adequate’ input maize-based agriculture in the derived savanna benchmark zone of Benin Republic. *Plant and Soil* 228:61–71.